

AD-A093 709

ARMY ELECTRONICS RESEARCH AND DEVELOPMENT COMMAND WS--ETC F/6 4/1
INITIAL FIELD MEASUREMENTS OF ATMOSPHERIC ABSORPTION AT 9 MICRO--ETC(U)
OCT 80 C W BRUCE, Y P YEE, B D HINDS

UNCLASSIFIED

ERADCOM/ASL-TR-0068


NL

100
100



1

END
DATE
FILMED
2-81
DTIC



AD A093709

ASL-TR-0068

LEVEL II

12

AD

Reports Control Symbol
OSD 1366

K

INITIAL FIELD MEASUREMENTS OF ATMOSPHERIC ABSORPTION AT 9 μ M TO 11 μ M WAVELENGTHS

OCTOBER 1980

By

C. W. BRUCE

Y. P. YEE

B. D. HINDS

R. G. PINNICK

US ARMY ATMOSPHERIC SCIENCES LABORATORY

White Sands Missile Range, NM 88002

R. J. BREWER

SCIENCE APPLICATIONS, INCORPORATED

Ann Arbor, MI

J. MINJARES

PHYSICAL SCIENCE LABORATORY

New Mexico State University
Las Cruces, NM 88003

DTIC
ELECTE
JAN 13 1981
E

Approved for public release; distribution unlimited

DNC FILE COPY



US Army Electronics Research and Development Command
ATMOSPHERIC SCIENCES LABORATORY
White Sands Missile Range, NM 88002

81 1 12 059

NOTICES

Disclaimers

The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

The citation of trade names and names of manufacturers in this report is not to be construed as official Government indorsement or approval of commercial products or services referenced herein.

Disposition

Destroy this report when it is no longer needed. Do not return it to the originator.

(14) ERADCOM/ASL-TR-0068

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ASL-TR-0068 /	2. GOVT ACCESSION NO. AD-A093709	3. RECIPIENT'S CATALOG NUMBER
6. TITLE (and Subtitle) INITIAL FIELD MEASUREMENTS OF ATMOSPHERIC ABSORPTION AT 9.2 TO 10.8 WAVELENGTHS.		4. TYPE OF REPORT & PERIOD COVERED Final Report.
7. AUTHOR(s) C. W. Bruce, Y. P. Yee, B. D. Hinds, and R. G. Pinnick, ASL; R. J. Brewer, SAI; J. Minjares, PSL		5. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS US Army Atmospheric Sciences Laboratory White Sands Missile Range, NM 88002		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Electronics Research and Development Command Adelphi, Maryland 20783		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS DA Task 1L161102B53A
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) Charles W. Bruce Young Paul Yee B. D. Hinds R. G. Pinnick Ralph J. Brewer		12. REPORT DATE October 1980
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		13. NUMBER OF PAGES 23
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
18. SUPPLEMENTARY NOTES		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Spectrophone CO ₂ laser Atmospheric particulates Dust measurements absorption coefficient Knollenberg light scattering aerosol measurements		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A field-adapted spectrophone system employing a tunable CO ₂ laser source (over wavelengths 9.2 to 10.8 μ m) was used to measure atmospheric gaseous and particulate absorption at an isolated desert southwestern location in the United States. Measurements were made both for ambient conditions (when aerosol particulate absorption was negligible compared to that of gases) and for dusty conditions resulting from vehicular traffic. For ambient conditions,		

FORM 1472 EDITION OF 1 NOV 65 IS OBSOLETE 1

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

410663

JOB

20. ABSTRACT (cont)

the gaseous absorption coefficient varied with time from expected levels upward by as much as a factor of 3. Sources which could be correlated with increased absorption are discussed. For dusty conditions the spectrophone data were compared with estimates of the absorption coefficient calculated on the basis of measured particle size distributions together with estimates of particle complex indices of refraction. Temporal variation of the absorption coefficient correlated closely for the two methods, while the calculated values were generally higher. Sampling and calculational uncertainties are suggested as likely to be responsible for this discrepancy.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	

SUMMARY

Measurements of both gaseous and aerosol atmospheric absorption made with a differential CO_2 laser spectrophone system have been reported. The spectrophone allows in situ measurements to be made on a real-time basis at frequencies of electro-optical systems and obviates some uncertainties present in other methods.

In the case of atmospheric gases, the use of this system shows how easily absorption may be significantly increased due to local and often innocuous appearing sources. Numerous trace gas pollutants at the 1 ppm level have absorption coefficients comparable to the ambient absorption and distinctive spectral patterns for the $9.2\mu\text{m}$ to $10.8\mu\text{m}$ emission line spectrum of a C^{17}O_2 laser. These factors suggest that the spectrophone device may be well-suited for air pollution studies.

Measurement of absorption by atmospheric particulates is more difficult than for gases. While the interaction between the spectrophone laser beam and the particles and the acoustical sensing are reasonably well understood, accurate sampling of all sizes of particles under windy conditions is difficult. (However, sampling particles with any other currently available technique faces similar difficulties.) One aspect of the importance of the particulate component is that of its extreme variability. Quiescent and low-wind conditions generally produce absorption coefficients that are negligible compared with the gaseous component at wavelengths around $10\mu\text{m}$ while high winds and/or anthropogenic disturbances can raise the particulate absorption by orders of magnitude well above the gaseous component.

ACKNOWLEDGMENT

The authors acknowledge the x-ray analysis performed by J. Whittler of the White Sands Missile Range Chemistry Laboratory and the aid of Mr. Glenn Hoidale of this laboratory for his many suggestions made during the preparation of this report.

CONTENTS

INTRODUCTION.....	7
BACKGROUND.....	7
MEASUREMENT SYSTEM.....	8
MEASUREMENTS AND ANALYSIS.....	13
REFERENCES.....	22
SELECTED BIBLIOGRAPHY.....	23

INTRODUCTION

Measurement of the absorption due to atmospheric gaseous and particulate matter at infrared wavelengths is a subject of current interest because of the relevance to communications, atmospheric sensing, and radiative transfer problems. Thus far, estimates of atmospheric absorption generally have been derived from measurements/estimates of the concentration of significant contributors at the wavelengths of interest. These concentration data are then converted to absorption coefficients at specific wavelengths. For gaseous absorption this may involve an insufficient data base of absorption coefficients, primarily for trace gases. In the case of particles, assumptions involving both shapes and complex indices of refraction must generally be made.¹

This report discusses the relatively direct in situ field measurements of both gaseous and particulate absorption made by using a CO₂ laser spectrophone system. The spectrophone technique obviates many of the problems associated with much less direct approaches currently used for obtaining atmospheric absorption. The measurements were made in the spring of 1978 at a remote desert site in the White Sands Missile Range (WSMR), New Mexico. Although both gaseous and particulate absorptions are discussed, the focus is on absorption by soil-derived dust generated by vehicular traffic.

BACKGROUND

In a previous paper² a spectrophone for in situ measurements of atmospheric gaseous and/or particulate absorption was described. Environmental chamber measurements on known atmospheric aerosols (quartz and calcite dust) were made with this system which allows continuous flow through sampling and a high degree of isolation from acoustic noise. Measurements for known atmospheric gaseous absorbers (methane,³ ammonia,⁴ and water vapor⁵) have also been

¹S. G. Jennings, R. G. Pinnick, and H. J. Auvermann, 1978, "Effects of Particulate Complex Refractive Index and Particle Size Distribution Variations on Atmospheric Extinction and Absorption for Visible through Middle-IR Wavelengths," Appl Opt, 17:3922-3928

²C. W. Bruce and R. G. Pinnick, 1977, "In-Situ Measurements of Aerosol Absorption with a Resonant CW Laser Spectrophone," J Appl Opt, 16:1762-1765

³C. W. Bruce et al, 1976, "Application of Pulsed-Source Spectrophone to Absorption by Methane at DF Laser Wavelengths," Appl Opt Letters, 15:2970-2972

⁴R. J. Brewer and C. W. Bruce, 1978, "Photoacoustic Spectroscopy of NH₃ at the 9 μ m and 10 μ m ¹²C ¹⁸O₂ Laser Wavelengths," J Appl Opt, 17:3746-3749

⁵K. O. White et al, 1978, "Water Vapor Continuum Absorption in the 3.5-4.0 μ m Region," Appl Opt, 17:2711-2720

made with similar spectrophones. In general, the gaseous absorption measurements were in good agreement with those obtained by using long path transmission cells with beam folding optics (White cells). In the case of aerosols, the measurements were compared to Mie theory predictions of absorption (for spherical homogeneous particles) based on measured particle size distributions and knowledge of particle complex refractive indices (which had previously been measured). Though the particles were quite irregular in shape, the measured and calculated results compared reasonably well (generally within a factor of 2) for the substances measured.

However, when atmospheric gases and dust are encountered in the field, as they were during this research, the presence of unknown trace gases and particles of undefined composition complicates any comparison of spectrophone measured absorption as compared with that predicted. For gases, comparison presumes distinctive (and well-known) spectral patterns at the probe laser wavelengths to permit identification of the species. The particulate comparison is more complex. In this case, both elements of the comparison are based on measurements which may involve sampling errors which are hard to define; for example, they are a function of windspeed and wind direction. Spectrophone operational parameters (chopping rates and power densities) must be chosen to ensure adequate response to the particles. Principal criteria for these choices were that the period of optical beam modulation exceed the thermal cooling time for the largest contributing absorbing particles and that probe beam power densities be low enough that optical properties not be significantly altered for contributing sizes of particles of a given complex index and thermal diffusivity. Since a light scattering counter was used to measure particle size and concentration, its response and size resolution for particles characteristic of soil-derived dust must be defined. Finally, dust composition must be determined. Additional complications arise from the fact that the dust composition can be a strong function of size⁴ (but unlikely in this case) and that refractive index information for the various dust constituents (which are required for Mie calculations of absorption) have been reported for only a few specific minerals.

MEASUREMENT SYSTEM

The ensemble of instrumentation consists of a spectrophone system for gaseous and particulate absorption measurements and correlative measurement instrumentation including a light scattering aerosol counter, a filter sampler, and a dew point hygrometer.

A schematic representation of the spectrophone system is shown in figure 1 where the laser beam path is traced through the alternate routes of spectrometer and spectrophones. Briefly, the differential resonant spectrophone system consists of two measurement elements. One samples both

⁴J. D. Lindberg and J. B. Gillespie, 1977, "Relationship Between Particle Size and Imaginary Refractive Index in Atmospheric Dust," Appl Opt., 16:2628-2630




Figure 1. Schematic (top view) of the portable optical table showing optical paths for both the CO₂ and the He-Ne alignment lasers, spectrometer for laser line identification, gas and total absorption spectrometers, power meter, and two of the auxiliary systems used, i.e., (EG&G) dew point hygrometer and dust sampling filter using 0.2 μ m Nuclepore filter elements. Flowmeters are for the spectrophones and the auxiliary measurements. BS and M refer to adjacent beam splitters or mirrors.

gaseous and particulate atmospheric contents through an unfiltered intake; a second, whose intake is filtered to remove particles, samples only the gaseous component. The gain-equalized difference signal from the two elements is due to the particles, while switching the total absorption leg of the difference signal to electrical ground gives only the molecular absorption.

Both spectrophones are cylindrically symmetric with the coaxial laser probe beam. The gas spectrophone has cell end windows, while the beam and atmospheric constituents enter the "total" (gas and particle) absorption spectrophone through the open upper end as illustrated in figure 2. Acoustic wave filters and a microphone designed to promote laminar flow occupy the central region, followed by a tube with relatively high acoustical inertance terminating in a calorimeter. This tube is a low pass filter that minimizes flow noise from the pump connection (where it is not yet laminar) and any signal due to the interaction of the laser beam with the calorimeter. The CO₂ laser used for these measurements is a modified commercial unit (GTE Sylvania model 948) tunable from approximately 9.15 μ m to 10.8 μ m in about 80 lines. Signal processing for the spectrophone system is indicated in the block diagram of figure 3. Pulsed laser sources could be substituted for the CW source and signal processing as described previously.¹

Considerable attention was given to the calibration of the light scattering counter (a Knollenberg model CSASP-100) used for measurement of aerosol size distributions and concentrations. The principle of operation of this instrument and determination of its response to spherical and nonspherical particles of various compositions are described in earlier papers.^{2,3} Briefly, the instrument works on the principle that as aerosol flows through an illuminated volume, light scattered by a single particle into a given (near forward scattering) solid angle is measured and used to determine particle size by electronically classifying response pulses according to their magnitude. To define the instrument's size resolution for measurement of soil-derived aerosols, we have measured its response to uniform irregular particles with refractive indices characteristic of those of soil dust constituents. The results, which are shown compared to theoretical response calculations for spheres of equal cross section in figure 4, show that the instrumental size resolution is defined by an envelope enclosing the theoretical response curves. This envelope (shown by the dashed curves in figure 4) indicates what uncertainty in particle size results from a certain response (or pulse height) measurement, considering the fact that the

¹C. W. Bruce et al, 1976, "Application of Pulsed-Source Spectrophone to Absorption by Methane at DF Laser Wavelengths," Appl Opt Letters, 15:2970-2972

²R. G. Pinnick and H. J. Auvermann, 1979, "Response Characteristics of Knollenberg Light-Scattering Aerosol Counters," J Aerosol Sci, 10:55-74

³R. G. Pinnick and J. M. Rosen, 1979, "Response of Knollenberg Light-Scattering Counters to Non-spherical Poublet Polystyrene Latex Aerosols," J Aerosol Sci, 10:533-538




Figure 2. Schematic diagram (side view) of the total (gas and particle) absorption spectrophone.




Figure 3. Signal processing for the CW laser resonant differential spectrophone system.




Figure 4. Knollenberg CSASP response: measured (circles) for nonspherical particles of doublet-shaped polystyrene (refractive index $m = 1.592-0i$), cubical sodium chloride ($m = 1.54-0i$), ellipsoidal potassium chlorate ($m = 1.409-0i$), and slightly nonspherical pollens and spores (puff balls, lycopodium powder, paper mulberry, ragweed, sweet vernal, and pecan) (with $m = 1.53-0i$); and calculated by using Mie theory (smooth solid curves) for spheres of equal cross section and refractive index. The envelope indicated by the smooth dashed curves defines an estimate of range of uncertainty in particle size that results for a particular response measurement for particles of unknown shape and refractive index.

particles are irregular and have different compositions (and refractive indices). This envelope has been extended slightly for application to soil dust constituents in an attempt to consider their slightly wider range of refractive indices and more irregular shapes. A curve drawn through the center of the envelope is then taken as the calibration for the instrument (to connect instrument response to particle size), and the envelope itself is used to define the particle sizing errors.

Air flows into both spectrophone and particle counter through intake bells which attempt to give relatively smooth flow through their measurement region and approximately isokinetic sampling when windspeeds are small.

MEASUREMENTS AND ANALYSIS

The CO₂ laser spectrophone system utilized for measurement of gaseous and particulate absorption generally can adequately define the spectral dependence of the particulate absorption in the 9 μ m to 11 μ m region, even though there are gaps between the four laser bands within this region. Molecular constituents cannot be completely characterized with the laser source since atmospheric gaseous absorption lines are more numerous than the laser probe lines and effectively narrower than the laser line spacing. However, the fixed patterns of the constituent gases generally do have strong, spectrally distinct contrasts for the absorbing constituents. This may permit identification of absorbing gaseous concentrations by deconvolution of the spectra using self-consistency.⁹

The measurements were performed partly to survey absorption by ambient atmospheric gases at the desert site. For purposes of gaseous analysis, measurements were made for 60 to 80 easily obtained laser lines with an estimated probable single measurement accuracy of 15 percent. However, significant changes in the absorption can occur in less time than is required for a complete manual spectral scan (about 1 to 2 hours). Such changes alter the observed spectral pattern, and deconvolution of the results may not give accurate concentrations by self-consistency of the spectra. Use of selected sets of laser probe lines is much more satisfactory since the measurement period can be made smaller than that for atmospheric changes. The technique used for choosing a minimum number of spectral lines is discussed by Samuel et al.⁹ Water vapor and carbon dioxide produce most of the ambient level atmospheric absorption in the 9 μ m to 11 μ m atmospheric transmission spectra "window." Absorption by atmospheric carbon dioxide measured with a C¹⁸O₂ laser probe is an exception to the assumption of distinctive spectra. Here the band profile of the absorption coefficient peaks varies relatively slow with wavelength. Examples of distinctive trace gas spectra are those of ozone, which appeared at levels as high as 0.030 ppm based on coefficients by

⁹C. Samuel, C. W. Bruce, and R. J. Brewer, 1978, Spectrophone Analysis of Gas Samples Obtained at Field Site, ASL-TR-0009, US Army Atmospheric Sciences Laboratory, White Sands Missile Range, NM

Patty,^{1*} and ammonia, which appeared at levels as high as 0.023 ppm based on coefficients by Brewer.²

For ambient atmospheric conditions, the spectrophone measured net absorption coefficients ranged from the expected ambient values to several times those values. The magnitude and temporal variation in the absorption coefficients during one 3-hour midday period for a number of CO₂ laser lines are shown in figure 5. As mentioned, these variations were larger than expected, as was the average level of the absorption. The major contribution was expected to stem from carbon dioxide at roughly 0.07 km⁻¹ and water vapor at about 0.04 km⁻¹ for the relative humidity encountered (about 3 torr). In previous samples of desert air analyzed in the laboratory, the 9 μ m to 11 μ m gaseous absorption coefficients were also often approximately twice the expected value.³ In those cases, possible contamination of the stainless sample system was suspected; however, the flowing sample changed in a matter of seconds, making contamination highly unlikely. Of course, the presence of personnel may have contributed to the high absorption. This potential problem will be avoided in the future by the development of an automated laser tuning system.

To point out the caution needed in interpreting the source of absorption in spectrophone measurements, several observations regarding interaction between personnel in the locale and the measurements might be of interest.

At one point, the absorption signal rose off scale (roughly 3x previous value) and persisted over several spectral lines. Then it was noticed that a meteorological observer about 15 m upwind was carrying a lighted cigarette. After the cigarette was extinguished, the absorption returned to previous levels.

On another occasion a particle counter was being cleaned with Freon 22 and acetone about 40 m downwind (windspeed about 7 m s⁻¹). The absorption coefficient increased when the cleaning began and decreased when the cleaning stopped. That the increase in absorption was due to these cleaning agents was obvious from the already familiar spectral dependence of these laboratory solvents. The increase (labeled t₂) relative to a more typical result (labeled t₁) is shown in the inset in figure 5 where a segment of the CO₂ laser 10 μ m P series is repeated for the two conditions.

^{1*}R. R. Patty et al, 1974, "CO₂ Laser Absorption Coefficients for Determining Ambient Levels of O₃, NH₃, and C₂H₆, Appl Opt, 13:2850-2854

²R. J. Brewer and C. W. Bruce, 1978, "Photoacoustic Spectroscopy of NH₃ at the 9 μ m and 10 μ m ¹²C O₂ Laser Wavelengths," J Appl Opt, 17:3746-3749

³C. Samuel, C. W. Bruce, and R. J. Brewer, 1978, Spectrophone Analysis of Gas Samples Obtained at Field Site, ASL-TR-0009, US Army Atmospheric Sciences Laboratory, White Sands Missile Range, NM




Figure 5. Atmospheric absorption coefficient data illustrating levels (points) and variations (vertical bar) for a 3-hour midday period (4 April 1978). Inset is a relative absorption (semilog) plot for 10μm P series CO₂ laser wavelength segment at two times whose significance is identified in the text. Straight lines connect measurement values for ease of identification.

A strong smell of creosote from a balloon enclosure persisted during some periods of high absorption, so an off-line spectrophone spectral analysis was performed in the laboratory on a sample taken from inside the enclosure. The results showed that creosote could not have been the source of the increase in absorption.

During measurements of atmospheric absorption under ambient conditions, the particulate absorption coefficient at $10\mu\text{m}$ was found to be generally less than 10 percent of the net gaseous absorption. Only occasional passage of vehicular traffic on the unpaved desert roads and one dust devil produced absorption coefficients larger than 10^{-1} km^{-1} .

Next, dust generated by vehicular traffic over the desert terrain was investigated. To correlate the spectrophone-measured absorption coefficient with an independent dust measurement, the (previously mentioned) Knollenberg probe was placed approximately 1 m from the spectrophone probe. The dust created by repeated circulation of a military truck vigorously driven over nearby desert terrain was then continuously measured with the spectrophone at a fixed CO_2 laser frequency while the counter accumulated data for consecutive 5-s intervals. The time resolution for the spectrophone was limited to about 1 s by the signal bandwidth used, and again the estimated probable error is 15 percent for particle radii less than $10\mu\text{m}$ (except for the elusive sampling errors).

Comparison of spectrophone measurements of absorption coefficient with calculations based on the Knollenberg data was then attempted. This comparison goes well beyond correlation of absorption coefficient with particle densities and requires some knowledge of the dust particle composition (or refractive index). Therefore, following the tests, vehicular dust which settled onto the ground was taken to the laboratory for spectral and chemical analysis.

The spectral analysis was accomplished by dispersing the sample dust in an environmental chamber and scanning the spectral region of concern using spectrophone systems similar to those used in the field. The reliability of these results depends to some degree on the assumption that the size distribution of redispersed dust samples is the same as it was in the field.¹

The chemical analysis was attempted by using two methods: x-ray diffractometry and infrared spectrophotometric analyses. The x-ray analysis revealed a strong quartz component but little else since not all of the dust consisted of crystalline material. The infrared spectrophotometric analysis was much more informative because of the distinct infrared spectral absorption

¹S. G. Jennings, R. G. Pinnick, and H. J. Auvermann, 1978, "Effects of Particulate Complex Refractive Index and Particle Size Distribution Variations on Atmospheric Extinction and Absorption for Visible through Middle-IR Wavelengths," Appl Opt, 17:3922-3928

patterns of the minerals involved.¹¹ In this analysis, the sample transmission spectra between wavelengths 2.5 μ m to 40 μ m were measured with a Perkin Elmer model 521 infrared grating spectrophotometer. The spectral transmission "fingerprints" of the substance under test were compared with those of known pure reference samples to assess the fractional composition of the unknown dust sample.¹² In this spectrometric estimate of mineral composition, the following assumptions were made: (1) the sample and reference particulate size spectra are identical functions of those distributions, (2) the absorption resonance peaks were assumed to be unique to a particular mineral, and (3) sample and reference minerals were assumed to be in the same chemical state. Under these assumptions the fractional mineral composition of the dust sample was then calculated by converting the transmission spectra to absorption coefficients and scaling the absorption resonance peaks linearly with mineral mass concentration. The results of this analysis are shown in table 1. It is noteworthy that the minerals identified (quartz, montmorillonite, calcite, and gypsum) were also found in naturally occurring aerosol samples collected within 4 km of where this test was conducted, although the proportions of each mineral were markedly different.¹¹

TABLE 1. RESULTS OF SOIL ANALYSIS WITH COMPLEX INDICES

Mineral	Identification Wavelength (μ m)	Fractional Composition (%)	Complex Index at 9.55 μ m (n_r n_i)
Gypsum	2.82	26	2.0, 0.30
Calcite	6.94	14	1.7, 1.20
Montmorillonite	9.80	35	0.86, 1.28
Quartz	12.82	25	1.0, 5.4

The fractional composition values of table 1 and the complex refractive indices of the constituents, with the Lorenz-Mie theory, can be used to

¹¹G. B. Hoidale and A. J. Blanco, 1969, "Infrared Absorption Spectra of Atmospheric Dust over an Interior Desert Basin," Pure and Appl Geophys. 74:151-164

¹²G. Duyckaerts, 1959, "The Infra-Red Analysis of Solid Substances," Analyst. 84:201-214

calculate the contributions to the total absorption coefficient of each component. Assumptions made in this process are those of particle sphericity, homogeneity, and similar size distributions for each constituent. In addition, for the birefringent quartz and calcite particles, the absorption was calculated by considering two populations of particles--one having refractive index of the ordinary ray and one the extraordinary ray--and adding the absorption coefficients for these fractional populations on a 2/3 to 1/3 basis.

Presently, even a rough estimate of the effect of these assumptions on the calculated absorption coefficients would be very difficult. Furthermore, the complex indices for these constituents are not well-known and montmorillonite (for example) does not have a fixed composition. Quartz is probably the best defined with respect to index; the others may be accurate within perhaps a factor of 2.

Figure 6 shows an example of the spectrophone measured dust absorption coefficients compared to those predicted from the Knollenberg counter dust size distribution measurements together with the spectrophotometric analysis of the dust composition. Knollenberg data were taken continuously during this 2-minute period, although the instrument particle count rate exceeded the maximum reliable rate (determined in the laboratory to be about 1000 s^{-1} under these conditions) for several 5-s accumulation intervals. (The predictions of absorption for this suspect data generally fall above the 100 km^{-1} level in figure 6.) In any case, the predicted absorption for all of the Knollenberg data displays similar temporal variation as the spectrophone measured absorption, although the values are higher by a factor of 3 to 5. The "error" bars shown for the Knollenberg-based predictions indicate only possible error caused by uncertainty in particle size determination made from the counter measurements; no estimate of errors accompanying the assumption of spherical particles, of the approximate treatment for birefringent particles, and of the assumption of similar size distribution for each dust mineral constituent was attempted.

Thus, in view of the multitude of rather critical assumptions made in predicting absorption from the Knollenberg size distribution data, and considering that the aerosol sampling losses may be significant and different for the spectrophone and Knollenberg sensors, agreement of the measured and predicted absorption in figure 6 is considered respectable.

To determine how much absorption (at the $9.21 \mu\text{m}$ wavelength) is contributed by particles of various sizes the differential absorption coefficients were calculated from the Knollenberg size distribution measurements. The results for 35 s of data (corresponding to times denoted a-g in figure 6) are presented in figure 7. These results suggest that the absorption is dominated by particles with radii in the $1 \mu\text{m}$ to $5 \mu\text{m}$ range regardless of dust loading.

Spectrophone measurements of the spectral dependence of the absorption for the vehicular dust redispersed in a laboratory environmental chamber are shown in figure 8. Because of the uncertainties mentioned earlier for the complex indices of the dust constituents, no attempt was made to calculate a result




Figure 6. Absorption coefficient of vehicle dust at the $9.21 \mu\text{m}$, R(34) CO_2 laser line as a function of time during the test: measured with in situ spectrophone (solid line); and calculated from particle size distribution measurements made with a Knollenberg light scattering aerosol counter (dashed line). The "error" bars superimposed on the Knollenberg-based predictions indicate only the error caused by uncertainty in particle size determination. Other sources of error in the predictions (see text) and differences in the spectrophone and Knollenberg probe sampling efficiencies are expected to cause the remaining discrepancy between measured and predicted absorption.




Figure 7. Differential absorption coefficients of vehicular dust showing that $1\mu\text{m}$ to $5\mu\text{m}$ (equivalent) radius particles dominate absorption at the $9.21\mu\text{m}$ CO_2 laser line. These coefficients are calculated from the Knollenberg size distribution measurements made at times a-g (in figure 6) together with estimates of the vehicular dust refractive indices (see text for additional details of this calculation).

for comparison. However, the spectral profiles for quartz and calcite dusts measured with spectrophones (for similar size distributions) and transmission spectra for montmorillonite and gypsum measured with the spectrophotometer techniques suggest that the strong absorption feature in the $9\mu\text{m}$ region in figure 8 is due partly to quartz, but that its broad character and its extension past $9.6\mu\text{m}$ is probably due to the presence of clay minerals such as montmorillonite.




Figure 8. Spectrophone absorption measurements of the spectral dependence of samples of vehicle dust.

REFERENCES

1. Jennings, S. G., R. G. Pinnick, and H. J. Auvermann, 1978, "Effects of Particulate Complex Refractive Index and Particle Size Distribution Variations on Atmospheric Extinction and Absorption for Visible through Middle-IR Wavelengths," Appl Opt, 17:3922-3928.
2. Bruce, C. W., and R. G. Pinnick, 1977, "In-Situ Measurements of Aerosol Absorption with a Resonant CW Laser Spectrophone," J Appl Opt, 16:1762-1765.
3. Bruce, C. W., et al, 1976, "Application of Pulsed-Source Spectrophone to Absorption by Methane at DF Laser Wavelengths," Appl Opt Letters, 15:2970-2972.
4. Brewer, R. J., and C. W. Bruce, 1978, "Photoacoustic Spectroscopy of NH_3 at the $9\mu\text{m}$ and $10\mu\text{m}$ $^{12}\text{C}^{18}\text{O}_2$ Laser Wavelengths," J Appl Opt, 17:3746-3749.
5. White, K. O., et al, 1978, "Water Vapor Continuum Absorption in the $3.5\text{--}4.0\mu\text{m}$ Region," Appl Opt, 17:2711-2720.
6. Lindberg, J. D., and J. B. Gillespie, 1977, "Relationship Between Particle Size and Imaginary Refractive Index in Atmospheric Dust," Appl Opt, 16:2628-2630.
7. Pinnick, R. G., and H. J. Auvermann, 1979, "Response Characteristics of Knollenberg Light-Scattering Aerosol Counters," J Aerosol Sci, 10:55-74.
8. Pinnick, R. G., and J. M. Rosen, 1979, "Response of Knollenberg Light-Scattering Counters to Non-spherical Poublet Polystyrene Latex Aerosols," J Aerosol Sci, 10:533-538.
9. Samuel, C., C. W. Bruce, and R. J. Brewer, 1978, Spectrophone Analysis of Gas Samples Obtained at Field Site, ASL-TR-0009, US Army Atmospheric Sciences Laboratory, White Sands Missile Range, NM.
10. Patty, R., G. M. Russwurm, W. A. McClenny, and D. R. Morgan, 1974, " CO_2 Laser Absorption Coefficients for Determining Ambient Levels of O_3 , NH_3 , and C_2H_2 ," Appl Opt, 13:2850-2854.
11. Hoidale, G. B., and A. J. Blanco, 1969, "Infrared Absorption Spectra of Atmospheric Dust over an Interior Desert Basin," Pure and Appl Geophys, 74:151-164.
12. Duyckaerts, G., 1959, "The Infra-Red Analysis of Solid Substances," Analyst, 84:201-214.

SELECTED BIBLIOGRAPHY

1. Spitzer, W. G., and D. A. Kleinman, 1961, "Infrared Lattice Sands of Quartz," Phys Rev, 121:1324-1335.
2. Toon, O. B., J. B. Pollack, and Carl Sagan, 1977, "Physical Properties of the Particles Composing the Martian Dust Storm of 1971-1972," Icarus, 30:663-696.

DISTRIBUTION LIST

Commander
US Army Aviation Center
ATTN: ATZQ-D-MA
Fort Rucker, AL 36362

Chief, Atmospheric Sciences Div
Code ES-81
NASA
Marshall Space Flight Center, AL 35812

Commander
US Army Missile Command
ATTN: DRDMI-RRR/Dr. O. M. Essenwanger
Redstone Arsenal, AL 35809

Commander
US Army Missile Command
ATTN: DRSMI-OG (B. W. Fowler)
Redstone Arsenal, AL 35809

Commander
US Army Missile R&D Command
ATTN: DRDMI-TEM (R. Haraway)
Redstone Arsenal, AL 35809

Redstone Scientific Information Center
ATTN: DRSMI-RPRD (Documents)
US Army Missile Command
Redstone Arsenal, AL 35809

Commander
HQ, Fort Huachuca
ATTN: Tech Ref Div
Fort Huachuca, AZ 85613

Commander
US Army Intelligence
Center & School
ATTN: ATSI-CD-MD
Fort Huachuca, AZ 85613

Commander
US Army Yuma Proving Ground
ATTN: Technical Library
Bldg 2105
Yuma, AZ 85364

Dr. Frank D. Eaton
Geophysical Institute
University of Alaska
Fairbanks, AK 99701

Naval Weapons Center
Code 3918
ATTN: Dr. A. Shlanta
China Lake, CA 93555

Commanding Officer
Naval Envir Prediction Rsch Facility
ATTN: Library
Monterey, CA 93940

Sylvania Elec Sys Western Div
ATTN: Technical Reports Lib
PO Box 205
Mountain View, CA 94040

Geophysics Officer
PMTC Code 3250
Pacific Missile Test Center
Point Mugu, CA 93042

Commander
Naval Ocean Systems Center
(Code 4473)
ATTN: Technical Library
San Diego, CA 92152

Meteorologist in Charge
Kwajalein Missile Range
PO Box 67
APO San Francisco, CA 96555

Director
NOAA/ERL/APCL R31
RB3-Room 567
Boulder, CO 80302

Dr. B. A. Silverman D-1200
Office of Atmos Resources Management
Water and Power Resources Service
PO Box 25007 Denver Federal Center, Bldg. 67
Denver, CO 80225

Hugh W. Albers (Executive Secretary)
CAO Subcommittee on Atmos Rsch
National Science Foundation Room 510
Washington, DC 2055

Dr. Eugene W. Bierly
Director, Division of Atmos Sciences
National Science Foundation
1800 G Street, N.W.
Washington, DC 20550

Commanding Officer
Naval Research Laboratory
Code 2627
Washington, DC 20375

Defense Communications Agency
Technical Library Center
Code 222
Washington, DC 20305

Director
Naval Research Laboratory
Code 5530
Washington, DC 20375

Dr. J. M. MacCallum
Naval Research Laboratory
Code 1409
Washington, DC 20375

HQDA (DAEN-RDM/Dr. de Percin)
Washington, DC 20314

The Library of Congress
ATTN: Exchange & Gift Div
Washington, DC 20540
2

Mil Asst for Atmos Sci Ofc of
the Undersecretary of Defense
for Rsch & Engr/E&LS - RM 3D129
The Pentagon
Washington, DC 20301

AFATL/DLODL
Technical Library
Eglin AFB, FL 32542

Naval Training Equipment Center
ATTN: Technical Information Center
Orlando, FL 32813

Technical Library
Chemical Systems Laboratory
Aberdeen Proving Ground, MD 21010

US Army Materiel Systems
Analysis Activity
ATTN: DRXS-MP
APG, MD 21005

Commander
ERADCOM
ATTN: DRDEL-PA/ILS/-ED
2800 Powder Mill Road
Adelphi, MD 20783

Commander
ERADCOM
ATTN: DRDEL-ST-T (Dr. B. Zarwyn)
2800 Powder Mill Road
Adelphi, MD 20783
02

Commander
Harry Diamond Laboratories
ATTN: DELHD-CO
2800 Powder Mill Road
Adelphi, MD 20783

Chief
Intel Mat Dev & Spt Ofc
ATTN: DELEW-WL-I
Bldg 4554
Fort George G. Mead, MD 20755

Acquisitions Section, IRDB-D823
Library & Info Svc Div, NOAA
6009 Executive Blvd.
Rockville, MD 20752

Naval Surface Weapons Center
White Oak Library
Silver Spring, MD 20910

Air Force Geophysics Laboratory
ATTN: LCC (A. S. Carten, Jr.)
Hanscom AFB, MA 01731

Air Force Geophysics Laboratory
ATTN: LYD
Hanscom AFB, MA 01731

Meteorology Division
AFGL/LY
Hanscom AFB, MA 01731

The Environmental Research
Institute of MI
ATTN: IRIA Library
PO Box 8618
Ann Arbor, MI 48107

Mr. William A. Main
USDA Forest Service
1407 S. Harrison Road
East Lansing, MI 48823

Dr. A. D. Belmont
Research Division
PO Box 1249
Control Data Corp
Minneapolis, MN 55440

Commander
Naval Oceanography Command
Bay St. Louis, MS 39529

Commanding Officer
US Army Armament R&D Command
ATTN: DRDAR-TSS Bldg 59
Dover, NJ 07801

Commander
ERADCOM Scientific Advisor
ATTN: DRDEL-SA
Fort Monmouth, NJ 07703

Commander
ERADCOM Tech Support Activity
ATTN: DELSD-L
Fort Monmouth, NJ 07703

Commander
HQ, US Army Avionics R&D Actv
ATTN: DAVAA-O
Fort Monmouth, NJ 07703

Commander
USA Elect Warfare Lab
ATTN: DELEW-DA (File Cy)
Fort Monmouth, NJ 07703

Commander
US Army Electronics R&D Command
ATTN: DELCS-S
Fort Monmouth, NJ 07703

Commander
US Army Satellite Comm Agency
ATTN: DRCPM-SC-3
Fort Monmouth, NJ 07703

Commander/Director
US Army Combat Survl & Target
Acquisition Laboratory
ATTN: DELCS-D
Fort Monmouth, NJ 07703

Director
Night Vision & Electro-Optics Laboratory
ATTN: DELNV-L (Dr. R. Buser)
Fort Belvoir, VA 22060

Project Manager, FIREFINDER
ATTN: DRCPM-FF
Fort Monmouth, NJ 07703

PM, Firefinder/REMBASS
ATTN: DRCPM-FFR-TM
Fort Monmouth, NJ 07703

6585 TG/WE
Holloman AFB, NM 88330

AFWL/Technical Library (SUL)
Kirtland AFB, NM 87117

AFWL/WE
Kirtland, AFB, NM 87117

TRASANA
ATTN: ATAA-SL (D. Anguiano)
WSMR, NM 88002

Commander
US Army White Sands Missile Range
ATTN: STEWS-PT-AL
White Sands Missile Range, NM 88002

Rome Air Development Center
ATTN: Documents Library
TSLD (Bette Smith)
Griffiss AFB, NY 13441

Environmental Protection Agency
Meteorology Laboratory, MD 80
Rsch Triangle Park, NC 27711

US Army Research Office
ATTN: DRXRO-PP
PO Box 12211
Rsch Triangle Park, NC 27709

Commandant
US Army Field Artillery School
ATTN: ATSF-CD-MS (Mr. Farmer)
Fort Sill, OK 73503

Commandant
US Army Field Artillery School
ATTN: ATSF-CF-R
Fort Sill, OK 73503

Commandant
US Army Field Artillery School
ATTN: Morris Swett Library
Fort Sill, OK 73503

Commander
US Army Dugway Proving Ground
ATTN: STEDP-MT-DA-M
(Mr. Paul Carlson)
Dugway, UT 84022

Commander
US Army Dugway Proving Ground
ATTN: MT-DA-L
Dugway, UT 84022

US Army Dugway Proving Ground
ATTN: STEDP-MT-DA-T
(Dr. W. A. Peterson)
Dugway, UT 84022

Inge Dirmhirn, Professor
Utah State University, UMC 48
Logan, UT 84322

Defense Technical Information Center
ATTN: DTIC-DDA-2
Cameron Station, Bldg. 5
Alexandria, VA 22314
12

Commanding Officer
US Army Foreign Sci & Tech Cen
ATTN: DRXST-IS1
220 7th Street, NE
Charlottesville, VA 22901

Naval Surface Weapons Center
Code G65
Dahlgren, VA 22448

Commander
US Army Night Vision
& Electro-Optics Lab
ATTN: DELNV-D
Fort Belvoir, VA 22060

Commander
USATRADOC
ATTN: ATCD-FA
Fort Monroe, VA 23651

Commander
USATRADOC
ATTN: ATCD-IR
Fort Monroe, VA 23651

Dept of the Air Force
5WW/DN
Langley AFB, VA 23665

US Army Nuclear & Cml Agency
ATTN: MONA-WE
Springfield, VA 22150

Director
US Army Signals Warfare Lab
ATTN: DELSW-OS (Dr. Burkhardt)
Vint Hill Farms Station
Warrenton, VA 22186

Commander
US Army Cold Regions Test Cen
ATTN: STECR-OP-PM
APO Seattle, WA 98733

ATMOSPHERIC SCIENCES RESEARCH PAPERS

1. Lindberg, J.D., "An Improvement to a Method for Measuring the Absorption Coefficient of Atmospheric Dust and other Strongly Absorbing Powders," ECOM-5565, July 1975.
2. Avara, Elton P., "Mesoscale Wind Shears Derived from Thermal Winds," ECOM-5566, July 1975.
3. Gómez, Richard B., and Joseph H. Pierluissi, "Incomplete Gamma Function Approximation for King's Strong-Line Transmittance Model," ECOM-5567, July 1975.
4. Blanco, A.J., and B.F. Engebos, "Ballistic Wind Weighting Functions for Tank Projectiles," ECOM-5568, August 1975.
5. Taylor, Fredrick J., Jack Smith, and Thomas H. Pries, "Crosswind Measurements through Pattern Recognition Techniques," ECOM-5569, July 1975.
6. Walters, D.L., "Crosswind Weighting Functions for Direct-Fire Projectiles," ECOM-5570, August 1975.
7. Duncan, Louis D., "An Improved Algorithm for the Iterated Minimal Information Solution for Remote Sounding of Temperature," ECOM-5571, August 1975.
8. Robbiani, Raymond L., "Tactical Field Demonstration of Mobile Weather Radar Set AN/TPS-41 at Fort Rucker, Alabama," ECOM-5572, August 1975.
9. Miers, B., G. Blackman, D. Langer, and N. Lorimier, "Analysis of SMS/GOES Film Data," ECOM-5573, September 1975.
10. Manquero, Carlos, Louis Duncan, and Rufus Bruce, "An Indication from Satellite Measurements of Atmospheric CO₂ Variability," ECOM-5574, September 1975.
11. Petracca, Carmine, and James D. Lindberg, "Installation and Operation of an Atmospheric Particulate Collector," ECOM-5575, September 1975.
12. Avara, Elton P., and George Alexander, "Empirical Investigation of Three Iterative Methods for Inverting the Radiative Transfer Equation," ECOM-5576, October 1975.
13. Alexander, George D., "A Digital Data Acquisition Interface for the SMS Direct Readout Ground Station — Concept and Preliminary Design," ECOM-5577, October 1975.
14. Cantor, Israel, "Enhancement of Point Source Thermal Radiation Under Clouds in a Nonattenuating Medium," ECOM-5578, October 1975.
15. Norton, Colburn, and Glenn Hoidale, "The Diurnal Variation of Mixing Height by Month over White Sands Missile Range, N.M.," ECOM-5579, November 1975.
16. Avara, Elton P., "On the Spectrum Analysis of Binary Data," ECOM-5580, November 1975.
17. Taylor, Fredrick J., Thomas H. Pries, and Chao-Huan Huang, "Optimal Wind Velocity Estimation," ECOM-5581, December 1975.
18. Avara, Elton P., "Some Effects of Autocorrelated and Cross-Correlated Noise on the Analysis of Variance," ECOM-5582, December 1975.
19. Gillespie, Patti S., R.L. Armstrong, and Kenneth O. White, "The Spectral Characteristics and Atmospheric CO₂ Absorption of the Ho³⁺YLF Laser at 2.05 μ m," ECOM-5583, December 1975.
20. Novlan, David J., "An Empirical Method of Forecasting Thunderstorms for the White Sands Missile Range," ECOM-5584, February 1976.
21. Avara, Elton P., "Randomization Effects in Hypothesis Testing with Autocorrelated Noise," ECOM-5585, February 1976.
22. Watkins, Wendell R., "Improvements in Long Path Absorption Cell Measurement," ECOM-5586, March 1976.
23. Thomas, Joe, George D. Alexander, and Marvin Dubbin, "SATTEL — An Army Dedicated Meteorological Telemetry System," ECOM-5587, March 1976.
24. Kennedy, Bruce W., and Delbert Bynum, "Army User Test Program for the RDT&E-XM-75 Meteorological Rocket," ECOM-5588, April 1976.

25. Barnett, Kenneth M., "A Description of the Artillery Meteorological Comparisons at White Sands Missile Range, October 1974 - December 1974 ('PASS' - Prototype Artillery [Meteorological] Subsystem)," ECOM-5589, April 1976.
26. Miller, Walter B., "Preliminary Analysis of Fall-of-Shot From Project 'PASS'," ECOM-5590, April 1976.
27. Avara, Elton P., "Error Analysis of Minimum Information and Smith's Direct Methods for Inverting the Radiative Transfer Equation," ECOM-5591, April 1976.
28. Yee, Young P., James D. Horn, and George Alexander, "Synoptic Thermal Wind Calculations from Radiosonde Observations Over the Southwestern United States," ECOM-5592, May 1976.
29. Duncan, Louis D., and Mary Ann Seagraves, "Applications of Empirical Corrections to NOAA-4 VTPR Observations," ECOM-5593, May 1976.
30. Miers, Bruce T., and Steve Weaver, "Applications of Meteorological Satellite Data to Weather Sensitive Army Operations," ECOM-5594, May 1976.
31. Sharenow, Moses, "Redesign and Improvement of Balloon ML-566," ECOM-5595, June, 1976.
32. Hansen, Frank V., "The Depth of the Surface Boundary Layer," ECOM-5596, June 1976.
33. Pinnick, R.G., and E.B. Stenmark, "Response Calculations for a Commercial Light-Scattering Aerosol Counter," ECOM-5597, July 1976.
34. Mason, J., and G.B. Hoidale, "Visibility as an Estimator of Infrared Transmittance," ECOM-5598, July 1976.
35. Bruce, Rufus E., Louis D. Duncan, and Joseph H. Pierluissi, "Experimental Study of the Relationship Between Radiosonde Temperatures and Radiometric-Area Temperatures," ECOM-5599, August 1976.
36. Duncan, Louis D., "Stratospheric Wind Shear Computed from Satellite Thermal Sounder Measurements," ECOM-5800, September 1976.
37. Taylor, F., P. Mohan, P. Joseph and T. Pries, "An All Digital Automated Wind Measurement System," ECOM-5801, September 1976.
38. Bruce, Charles, "Development of Spectrophones for CW and Pulsed Radiation Sources," ECOM-5802, September 1976.
39. Duncan, Louis D., and Mary Ann Seagraves, "Another Method for Estimating Clear Column Radiances," ECOM-5803, October 1976.
40. Blanco, Abel J., and Larry E. Taylor, "Artillery Meteorological Analysis of Project Pass," ECOM-5804, October 1976.
41. Miller, Walter, and Bernard Engebos, "A Mathematical Structure for Refinement of Sound Ranging Estimates," ECOM-5805, November, 1976.
42. Gillespie, James B., and James D. Lindberg, "A Method to Obtain Diffuse Reflectance Measurements from 1.0 to 3.0 μm Using a Cary 171 Spectrophotometer," ECOM-5806, November 1976.
43. Rubio, Roberto, and Robert O. Olsen, "A Study of the Effects of Temperature Variations on Radio Wave Absorption," ECOM-5807, November 1976.
44. Ballard, Harold N., "Temperature Measurements in the Stratosphere from Balloon-Borne Instrument Platforms, 1968-1975," ECOM-5808, December 1976.
45. Monahan, H.H., "An Approach to the Short-Range Prediction of Early Morning Radiation Fog," ECOM-5809, January 1977.
46. Engebos, Bernard Francis, "Introduction to Multiple State Multiple Action Decision Theory and Its Relation to Mixing Structures," ECOM-5810, January 1977.
47. Low, Richard D.H., "Effects of Cloud Particles on Remote Sensing from Space in the 10-Micrometer Infrared Region," ECOM-5811, January 1977.
48. Bonner, Robert S., and R. Newton, "Application of the AN/GVS-5 Laser Rangefinder to Cloud Base Height Measurements," ECOM-5812, February 1977.
49. Rubio, Roberto, "Lidar Detection of Subvisible Reentry Vehicle Erosive Atmospheric Material," ECOM-5813, March 1977.
50. Low, Richard D.H., and J.D. Horn, "Mesoscale Determination of Cloud-Top Height: Problems and Solutions," ECOM-5814, March 1977.

51. Duncan, Louis D., and Mary Ann Seagraves, "Evaluation of the NOAA-4 VTPR Thermal Winds for Nuclear Fallout Predictions," ECOM-5815, March 1977.
52. Randhawa, Jagir S., M. Izquierdo, Carlos McDonald and Zvi Salpeter, "Stratospheric Ozone Density as Measured by a Chemiluminescent Sensor During the Stratcom VI-A Flight," ECOM-5816, April 1977.
53. Rubio, Roberto, and Mike Izquierdo, "Measurements of Net Atmospheric Irradiance in the 0.7- to 2.8-Micrometer Infrared Region," ECOM-5817, May 1977.
54. Ballard, Harold N., Jose M. Serna, and Frank P. Hudson Consultant for Chemical Kinetics, "Calculation of Selected Atmospheric Composition Parameters for the Mid-Latitude, September Stratosphere," ECOM-5818, May 1977.
55. Mitchell, J.D., R.S. Sagar, and R.O. Olsen, "Positive Ions in the Middle Atmosphere During Sunrise Conditions," ECOM-5819, May 1977.
56. White, Kenneth O., Wendell R. Watkins, Stuart A. Schleusener, and Ronald L. Johnson, "Solid-State Laser Wavelength Identification Using a Reference Absorber," ECOM-5820, June 1977.
57. Watkins, Wendell R., and Richard G. Dixon, "Automation of Long-Path Absorption Cell Measurements," ECOM-5821, June 1977.
58. Taylor, S.E., J.M. Davis, and J.B. Mason, "Analysis of Observed Soil Skin Moisture Effects on Reflectance," ECOM-5822, June 1977.
59. Duncan, Louis D. and Mary Ann Seagraves, "Fallout Predictions Computed from Satellite Derived Winds," ECOM-5823, June 1977.
60. Snider, D.E., D.G. Murcray, F.H. Murcray, and W.J. Williams, "Investigation of High-Altitude Enhanced Infrared Background Emissions" (U), SECRET, ECOM-5824, June 1977.
61. Dubbin, Marvin H. and Dennis Hall, "Synchronous Meteorological Satellite Direct Readout Ground System Digital Video Electronics," ECOM-5825, June 1977.
62. Miller, W., and B. Engebos, "A Preliminary Analysis of Two Sound Ranging Algorithms," ECOM-5826, July 1977.
63. Kennedy, Bruce W., and James K. Luers, "Ballistic Sphere Techniques for Measuring Atmospheric Parameters," ECOM-5827, July 1977.
64. Duncan, Louis D., "Zenith Angle Variation of Satellite Thermal Sounder Measurements," ECOM-5828, August 1977.
65. Hansen, Frank V., "The Critical Richardson Number," ECOM-5829, September 1977.
66. Ballard, Harold N., and Frank P. Hudson (Compilers), "Stratospheric Composition Balloon-Borne Experiment," ECOM-5830, October 1977.
67. Barr, William C., and Arnold C. Peterson, "Wind Measuring Accuracy Test of Meteorological Systems," ECOM-5831, November 1977.
68. Ethridge, G.A. and F.V. Hansen, "Atmospheric Diffusion: Similarity Theory and Empirical Derivations for Use in Boundary Layer Diffusion Problems," ECOM-5832, November 1977.
69. Low, Richard D.H., "The Internal Cloud Radiation Field and a Technique for Determining Cloud Blackness," ECOM-5833, December 1977.
70. Watkins, Wendell R., Kenneth O. White, Charles W. Bruce, Donald L. Walters, and James D. Lindberg, "Measurements Required for Prediction of High Energy Laser Transmission," ECOM-5834, December 1977.
71. Rubio, Robert, "Investigation of Abrupt Decreases in Atmospherically Backscattered Laser Energy," ECOM-5835, December 1977.
72. Monahan, H.H. and R.M. Cionco, "An Interpretative Review of Existing Capabilities for Measuring and Forecasting Selected Weather Variables (Emphasizing Remote Means)," ASL-TR-0001, January 1978.
73. Heaps, Melvin G., "The 1979 Solar Eclipse and Validation of D-Region Models," ASL-TR-0002, March 1978.

74. Jennings, S.G., and J.B. Gillespie, "M.I.E. Theory Sensitivity Studies - The Effects of Aerosol Complex Refractive Index and Size Distribution Variations on Extinction and Absorption Coefficients Part II: Analysis of the Computational Results," ASL-TR-0003, March 1978.
75. White, Kenneth O. et al, "Water Vapor Continuum Absorption in the 3.5 μ m to 4.0 μ m Region," ASL-TR-0004, March 1978.
76. Olsen, Robert O., and Bruce W. Kennedy, "ABRES Pretest Atmospheric Measurements," ASL-TR-0005, April 1978.
77. Ballard, Harold N., Jose M. Serna, and Frank P. Hudson, "Calculation of Atmospheric Composition in the High Latitude September Stratosphere," ASL-TR-0006, May 1978.
78. Watkins, Wendell R. et al, "Water Vapor Absorption Coefficients at HF Laser Wavelengths," ASL-TR-0007, May 1978.
79. Hansen, Frank V., "The Growth and Prediction of Nocturnal Inversions," ASL-TR-0008, May 1978.
80. Samuel, Christine, Charles Bruce, and Ralph Brewer, "Spectrophone Analysis of Gas Samples Obtained at Field Site," ASL-TR-0009, June 1978.
81. Pinnick, R.G. et al., "Vertical Structure in Atmospheric Fog and Haze and its Effects on IR Extinction," ASL-TR-0010, July 1978.
82. Low, Richard D.H., Louis D. Duncan, and Richard B. Gomez, "The Microphysical Basis of Fog Optical Characterization," ASL-TR-0011, August 1978.
83. Heaps, Melvin G., "The Effect of a Solar Proton Event on the Minor Neutral Constituents of the Summer Polar Mesosphere," ASL-TR-0012, August 1978.
84. Mason, James B., "Light Attenuation in Falling Snow," ASL-TR-0013, August 1978.
85. Blanco, Abel J., "Long-Range Artillery Sound Ranging: "PASS" Meteorological Application," ASL-TR-0014, September 1978.
86. Heaps, M.G., and F.E. Niles, "Modeling the Ion Chemistry of the D-Region: A case Study Based Upon the 1966 Total Solar Eclipse," ASL-TR-0015, September 1978.
87. Jennings, S.G., and R.G. Pinnick, "Effects of Particulate Complex Refractive Index and Particle Size Distribution Variations on Atmospheric Extinction and Absorption for Visible Through Middle-Infrared Wavelengths," ASL-TR-0016, September 1978.
88. Watkins, Wendell R., Kenneth O. White, Lanny R. Bower, and Brian Z. Sojka, "Pressure Dependence of the Water Vapor Continuum Absorption in the 3.5- to 4.0-Micrometer Region," ASL-TR-0017, September 1978.
89. Miller, W.B., and B.F. Engebos, "Behavior of Four Sound Ranging Techniques in an Idealized Physical Environment," ASL-TR-0018, September 1978.
90. Gomez, Richard G., "Effectiveness Studies of the CBU-88/B Bomb, Cluster, Smoke Weapon" (U), CONFIDENTIAL ASL-TR-0019, September 1978.
91. Miller, August, Richard C. Shirkey, and Mary Ann Seagraves, "Calculation of Thermal Emission from Aerosols Using the Doubling Technique," ASL-TR-0020, November, 1978.
92. Lindberg, James D. et al., "Measured Effects of Battlefield Dust and Smoke on Visible, Infrared, and Millimeter Wavelengths Propagation: A Preliminary Report on Dusty Infrared Test-I (DIRT-I)," ASL-TR-0021, January 1979.
93. Kennedy, Bruce W., Arthur Kinghorn, and B.R. Hixon, "Engineering Flight Tests of Range Meteorological Sounding System Radiosonde," ASL-TR-0022, February 1979.
94. Rubio, Roberto, and Don Hooch, "Microwave Effective Earth Radius Factor Variability at Wiesbaden and Balboa," ASL-TR-0023, February 1979.
95. Low, Richard D.H., "A Theoretical Investigation of Cloud/Fog Optical Properties and Their Spectral Correlations," ASL-TR-0024, February 1979.

96. Pinnick, R.G., and H.J. Auvermann, "Response Characteristics of Knollenberg Light-Scattering Aerosol Counters," ASL-TR-0025, February 1979.
97. Heaps, Melvin G., Robert O. Olsen, and Warren W. Berning, "Solar Eclipse 1979, Atmospheric Sciences Laboratory Program Overview," ASL-TR-0026 February 1979.
98. Blanco, Abel J., "Long-Range Artillery Sound Ranging: 'PASS' GR-8 Sound Ranging Data," ASL-TR-0027, March 1979.
99. Kennedy, Bruce W., and Jose M. Serna, "Meteorological Rocket Network System Reliability," ASL-TR-0028, March 1979.
100. Swingle, Donald M., "Effects of Arrival Time Errors in Weighted Range Equation Solutions for Linear Base Sound Ranging," ASL-TR-0029, April 1979.
101. Umstead, Robert K., Ricardo Pena, and Frank V. Hansen, "KWIK: An Algorithm for Calculating Munition Expenditures for Smoke Screening/Obscuration in Tactical Situations," ASL-TR-0030, April 1979.
102. D'Arcy, Edward M., "Accuracy Validation of the Modified Nike Hercules Radar," ASL-TR-0031, May 1979.
103. Rodriguez, Ruben, "Evaluation of the Passive Remote Crosswind Sensor," ASL-TR-0032, May 1979.
104. Barber, T.L., and R. Rodriguez, "Transit Time Lidar Measurement of Near-Surface Winds in the Atmosphere," ASL-TR-0033, May 1979.
105. Low, Richard D.H., Louis D. Duncan, and Y.Y. Roger R. Hsiao, "Microphysical and Optical Properties of California Coastal Fogs at Fort Ord," ASL-TR-0034, June 1979.
106. Rodriguez, Ruben, and William J. Vechione, "Evaluation of the Saturation Resistant Crosswind Sensor," ASL-TR-0035, July 1979.
107. Ohmstede, William D., "The Dynamics of Material Layers," ASL-TR-0036, July 1979.
108. Pinnick, R.G., S.G. Jennings, Petr Chylek, and H.J. Auvermann "Relationships between IR Extinction, Absorption, and Liquid Water Content of Fogs," ASL-TR-0037, August 1979.
109. Rodriguez, Ruben, and William J. Vechione, "Performance Evaluation of the Optical Crosswind Profiler," ASL-TR-0038, August 1979.
110. Miers, Bruce T., "Precipitation Estimation Using Satellite Data" ASL-TR-0039, September 1979.
111. Dickson, David H., and Charles M. Sonnenschein, "Helicopter Remote Wind Sensor System Description," ASL-TR-0040, September 1979.
112. Heaps, Melvin, G., and Joseph M. Heimerl, "Validation of the Dairchem Code, I: Quiet Midlatitude Conditions," ASL-TR-0041, September 1979.
113. Bonner, Robert S., and William J. Lentz, "The Visioceilometer: A Portable Cloud Height and Visibility Indicator," ASL-TR-0042, October 1979.
114. Cohn, Stephen L., "The Role of Atmospheric Sulfates in Battlefield Obscurations," ASL-TR-0043, October 1979.
115. Fawbush, E.J. et al, "Characterization of Atmospheric Conditions at the High Energy Laser System Test Facility (HELSTF), White Sands Missile Range, New Mexico, Part I, 24 March to 8 April 1977," ASL-TR-0044, November 1979.
116. Barber, Ted L., "Short-Time Mass Variation in Natural Atmospheric Dust," ASL-TR-0045, November 1979.
117. Low, Richard D.H., "Fog Evolution in the Visible and Infrared Spectral Regions and its Meaning in Optical Modeling," ASL-TR-0046, December 1979.
118. Duncan, Louis D. et al, "The Electro-Optical Systems Atmospheric Effects Library, Volume I: Technical Documentation, ASL-TR-0047, December 1979.
119. Shirkey, R. C. et al, "Interim E-O SAEL, Volume II, Users Manual," ASL-TR-0048, December 1979.
120. Kobayashi, H.K., "Atmospheric Effects on Millimeter Radio Waves," ASL-TR-0049, January 1980.
121. Seagraves, Mary Ann and Duncan, Louis D., "An Analysis of Transmittances Measured Through Battlefield Dust Clouds," ASL-TR-0060, February, 1980.

122. Dickson, David H., and Jon E. Ottesen, "Helicopter Remote Wind Sensor Flight Test," ASL-TR-0051, February 1980.
123. Pinnick, R. G., and S. G. Jennings, "Relationships Between Radiative Properties and Mass Content of Phosphoric Acid, HC, Petroleum Oil, and Sulfuric Acid Military Smokes," ASL-TR-0052, April 1980.
124. Hinds, B. D., and J. B. Gillespie, "Optical Characterization of Atmospheric Particulates on San Nicolas Island, California," ASL-TR-0053, April 1980.
125. Miers, Bruce T., "Precipitation Estimation for Military Hydrology," ASL-TR-0054, April 1980.
126. Stenmark, Ernest B., "Objective Quality Control of Artillery Computer Meteorological Messages," ASL-TR-0055, April 1980.
127. Duncan, Louis D., and Richard D. H. Low, "Bimodal Size Distribution Models for Fogs at Meppen, Germany," ASL-TR-0056, April 1980.
128. Olsen, Robert O., and Jagir S. Randhawa, "The Influence of Atmospheric Dynamics on Ozone and Temperature Structure," ASL-TR-0057, May 1980.
129. Kennedy, Bruce W., et al, "Dusty Infrared Test-II (DIRT-II) Program," ASL-TR-0058, May 1980.
130. Heaps, Melvin G., Roberts O. Olsen, Warren Berning, John Cross, and Arthur Gilcrease, "1979 Solar Eclipse, Part I - Atmospheric Sciences Laboratory Field Program Summary," ASL-TR-0059, May 1980.
131. Miller, Walter B., "User's Guide for Passive Target Acquisition Program Two (PTAP-2)," ASL-TR-0060, June 1980.
132. Holt, E. H., H. H. Monahan, and E. J. Fawbush, "Atmospheric Data Requirements for Battlefield Obscuration Applications," ASL-TR-0061, June 1980.
133. Shirkey, Richard C., August Miller, George H. Goedecke, and Yugal Behl, "Single Scattering Code AGAUSX: Theory, Applications, Comparisons, and Listing," ASL-TR-0062, July 1980.
134. Sojka, Brain Z., and Kenneth O. White, "Evaluation of Specialized Photoacoustic Absorption Chambers for Near-millimeter Wave (NMMW) Propagation Measurements," ASL-TR-0063, August 1980.
135. Bruce, Charles W., Young Paul Yee, and S. G. Jennings, "In Situ Measurement of the Ratio of Aerosol Absorption to Extinction Coefficient," ASL-TR-0064, August 1980.
136. Yee, Young Paul, Charles W. Bruce, and Ralph J. Brewer, "Gaseous/Particulate Absorption Studies at WSMR using Laser Sourced Spectrophones," ASL-TR-0065, June 1980.
137. Lindberg, James D., Radon B. Loveland, Melvin Heaps, James B. Gillespie, and Andrew F. Lewis, "Battlefield Dust and Atmospheric Characterization Measurements During West German Summertime Conditions in Support of Grafenwohr Tests," ASL-TR-0066, September 1980.
138. Vechione, W. J., "Evaluation of the Environmental Instruments, Incorporated Series 200 Dual Component Wind Set," ASL-TR-0067, September 1980.
139. Bruce, C. W., Y. P. Yee, B. D. Hinds, R. G. Pinnick, R. J. Brewer, and J. Minjares, "Initial Field Measurements of Atmospheric Absorption at 9 μ m to 11 μ m Wavelengths," ASL-TR-0068, October 1980.